



ZOOLOGICA

SCIENTIFIC CONTRIBUTIONS OF THE NEW YORK ZOOLOGICAL SOCIETY

FROM THE TROPICAL RESEARCH
STATION IN BRITISH GUIANA



VOLUME III. NUMBER 6

(Tropical Research Station Contribution Number 100)


DESCRIPTIONS OF NEW SPECIES OF COLEOPTERA

By E. A. SCHWARZ and H. S. BARBER

United States Department of Agriculture

PUBLISHED BY THE SOCIETY
THE ZOOLOGICAL PARK, NEW YORK

DECEMBER 24, 1921



DESCRIPTIONS OF NEW SPECIES OF COLEOPTERA

By E. A. SCHWARZ and H. S. BARBER

United States Department of Agriculture

Specimens of three species of beetles from Kartabo, Bartica District, British Guiana, have been submitted to us by their discoverer, Dr. W. M. Wheeler, for identification. Two of the species belong to the family Silvanidæ, while the third is an inconspicuous species of the ubiquitous Coccinellid genus *Scymnus*. The two first are, however, very interesting forms, and it has been necessary to erect a new genus to receive the larger and more abundant species. These, Dr. Wheeler informs us, were occupying the petioles of a tree (*Tachigalia*).

Coccidotrophus gen. nov. (Silvanidæ).

Eyes small, not emarginate, not prominent, occupying middle fourth of side of head. Antennæ inserted laterally, in deep cavity between frontal margin and prominent gena, close to and equidistant from base of mandible and front margin of eye; compactly eleven-jointed, club not abruptly enlarged, all joints traverse except basal joint; joints 3-9 rapidly increasing in size, the three-jointed club densely covered with fine short hairs. Antennal grooves short, distinct, parallel, bordered by lower edge of eye and carina from gena. Mouthparts free, maxillary palpi four-jointed, labial palpi three-jointed. Elytra with lateral costa and nine distinct series of striae punctures. Pygidium usually covered by elytra when abdomen is contracted (when distended the pygidium is often exposed). Intercoxal process of prosternum about one-seventh as wide as prothorax, wider behind, its sides overlapping the coxal cavities; front coxae separated from hind margin of prothorax by about their own length. Intercoxal process of mesosternum slightly narrower than that of prosternum, truncate apically and nearly reaching hind margin of coxae, metepimeron reaching the middle coxae. Hind coxae separated by strong,

acute process of first abdominal sternite, which fits into median notch in metasternum. First ventral abdominal segment with coxal lines very fine, short, curved, becoming almost parallel behind, but not reaching posterior margin; four following sternites each slightly shorter than the first, the last rounded apically, concave beneath, with carinate posterior margin. Legs short, stout, femora incrassate, first three tarsal joints inflated, hairy beneath, fourth small but distinct, fifth joint as long as the three basal joints together. Genotype: *Coccidotrophus socialis* sp. nov.

This genus is so unlike other Silvanids known to us that its true affinity was not recognized until Dr. Böving's studies of its larva proved it to agree well with typical Silvanid larvæ and not to be related to *Hapalips*, with which we were trying to associate the adults. Once suggested this relationship was readily confirmed by the characters used by Ganglbauer (*Die Käfer von Mitteleuropa*, 1889, vol. 3, p. 577) to distinguish the adults of this family: front coxal cavities closed behind, metepimeron reaching middle coxae, etc.

We cannot identify *Coccidotrophus* with any of the genera considered by Grouvelle (*Ann. Soc. Ent. Fr.*, vol. 81, 1912, pp. 313-386), but it seems most nearly to approach *Synoemis* Pascoe (1866), the only species of which was described from the Malayan Peninsula.

***Coccidotrophus socialis* sp. nov.**

(Plate VI, figs. 1 to 5).

Very elongate, parallel, depressed, shining, castaneous, glabrous (except for sparse, microscopical, decumbent pubescence).

Length, 3.5-4.5; width, 0.6-0.8 mm.

Habitat—British Guiana.

Head slightly wider than prothorax, widest at the very slightly prominent eyes, one-sixth longer than wide, feebly constricted into a neck in basal fourth, sides convergent in front, front margin broadly, shallowly notched; upper surface finely granulose except a small occipital smooth area, feebly convex, transversely somewhat tumid between and behind eyes, which

are situated about middle of sides; clypeus not separated from front, produced into a strong marginal carina, surface convex medially, concave laterally, especially over antennal sockets; labrum very short, transverse, feebly chitinized, emarginate in front and with a row of about six stiff hairs; gular region strongly pilose, feebly concave; mandibles moderately prominent, strongly bidentate apically; antennæ stout, shorter than width of front at their point of insertion. Pronotum three-fourths as wide as long, widest at apical angles which are subacute; narrowest just before middle and at base; sides feebly sinuate, subparallel, front margin straight, hind angles obtuse, hind margin arcuate, surface longitudinally flat, transversely feebly convex, sparsely punctate, the punctures elongate, side margins with fine marginal line. Scutellum transverse, widest behind, finely punctate. Elytra slightly wider than pronotum, more than three times as long as wide, base emarginate, humeral angles subacute, sides subparallel to apical fourth, thence conjointly rounded; surface longitudinally somewhat convex, transversely rather strongly convex, stria punctures moderate, the interstices each with a series of slightly smaller punctures supporting fine, decumbent, short hairs. Under surface shining but with irregular microscopic sculpture between the sparsely scattered, nearly obsolete punctures which bear the short, fine decumbent hairs.

Described from thirty-two examples from a large series collected by Dr. Wm. M. Wheeler at Kartabo in July and August, 1920. The sexes are almost indistinguishable unless the "palps" of the ovipositor are extruded.

Characters of legs, antennæ and mouth are shown in accompanying figures. Type, allotype and paratypes.—Cat. No. 24070, U.S.N.M.

Eunausibius Grouvelle, 1912.

Although we have seen neither of the two species (*Nausibius tenebrionides* and *N. elongatus* Grouv.) upon which Grouvelle, (Ann. Soc. Ent. Fr., vol. 81, 1912, p. 314) established this genus, his generic diagnosis applies so well to the species here described that his genus is adopted. The postcoxal lines are very feeble and difficult to see but are bent abruptly forward to the coxal cavity enclosing a small area under the trochanter:

the antennal club seems to be more abrupt and larger, and the shape of the produced clypeus differs also from the figures Grouvelle, (Ann. Soc. Ent. Fr., vol. 65, 1896, p. 193) has given of the two previously known species.

Eunausibius wheeleri sp. nov.

(Plate VI, figs. 6 to 10).

Elongate, parallel, moderately convex, smooth, shining, glabrous, pale castaneous.

Length, 3.0-3.5 mm.; width, 0.6-0.75 mm.

Habitat.—British Guiana.

Head wider than prothorax; eyes rather large and prominent, coarsely granulated, front widest in front of the rather strongly impressed fovea opposite which the margin is somewhat thickened; front margin broadly, feebly emarginate, front angles obliquely truncate; surface finely granulate, feebly convex. Labrum membranous, concealed beneath front. Gular area feebly concave, opaque, finely pubescent, the lateral carinae bordering the antennal grooves convergent posteriorly and passing middle of eye. Antennae as long as width of clypeus, club abruptly widened, oval. Pronotum about three-fourths as wide as long, widest at the slightly acute front angles, sides almost straight and parallel, finely carinate; front margin straight except small sinuation near angles, hind margin arcuate at middle, on each side straight, hind angles obtuse. Surface transversely convex, longitudinally flat except for a pair of feeble impressions at basal fourth. Scutellum transversely oval, twice as wide as long. Elytra slightly wider than prothorax, three times as long as wide, sides parallel, apices evenly rounded; surface nearly smooth, striae punctures feebly impressed but conspicuous by coloration below surface. Callow specimens display interstrial rows of microscopic appressed hairs. Under surface of body sparsely clothed with microscopic decumbent hairs, each set in a broad obsolescent puncture.

Described from eighteen specimens submitted by Dr. Wheeler, to whom the species is dedicated. Two of the specimens, dissected and mounted on slides, are males. One specimen, the allotype, is somewhat crushed and displays female sex

organs. We are unable to distinguish the sex of the other fifteen paratypes.

Type, allotype and paratypes.—Cat. No. 24071, U. S. N. M.
The antennæ and legs are shown in accompanying figures.

Scymnus Kugelann 1794 (Coccinellidæ)

After some hesitation we have decided to offer the following description of what appears to be a new but very commonplace species of this genus, since we have failed to find any description applicable thereto. It will, perhaps, be long before the multitude of tropical species of *Scymnus* will be determinable.

Scymnus (Diomus) xantholeucus sp. nov.

Oval, very convex, shining, pubescent, pale yellow except basal two-thirds of elytra, meso- and metasternum and median third of first two abdominal segments, which are infuscate.

Length, 1.7 mm.; width, 1.2 mm.

Habitat.—British Guiana.

Head and pronotum finely, rather densely punctate, elytra slightly more coarsely punctured; pubescence rather dense, the short, silky, suberect hairs bent in all directions forming no pattern. Scutellum pale. Elytra piceous in basal two-thirds, apically flavescens, the pale area not sharply limited. Under surface moderately, densely and finely punctate, the pubescence decumbent and regular. Prosternum with carinae convergent anteriorly, reaching front margin and uniting in an arc; the enclosed area feebly concave and supporting moderately long hairs. Post-metacoxal line as in other species of *Diomus* (Group A of Horn 1895). Penultimate abdominal sternite very broadly and feebly emarginate in the male.

Type (male), allotype and paratypes.—Cat. No. 24084, U.S.N.M.

Described from two males and two females reared by Dr. Wheeler from larvae found among coccids in company with the two species of *Silvanidæ*.

Although not related to *semiruber* Casey, the foregoing species is similar in shape and plan of coloration. A prescutellar infuscate area is suggested in one example.




PLATE VI.

- Fig. 1. *Coccidotrophus socialis* sp. nov. Right side of head, with antenna and mouth parts, ventral view. X 53.5.
 Fig. 2. Fore leg of same. X 36.5.
 Fig. 3. Middle leg of same. X 36.5.
 Fig. 4. Hind leg of same. X 36.5.
 Fig. 5. Hind wing of same. X 36.5.
 Fig. 6. *Eunausibius wheeleri* sp. nov. Antenna. X 58.5.
 Fig. 7. Fore leg of same. X 36.5.
 Fig. 8. Middle leg of same. X 36.5.
 Fig. 9. Hind leg of same. X 36.5.
 Fig. 10. Hind wing of same. X 36.5.

QL Wheeler, William Morton.
591 Study of some social
B74W48 beetles in British Guiana,
Ent. 1921.

